

### Motrin® Dosing for Infants and Children

**mL** = milliliter  
**tsp** = teaspoonful

If possible, use weight to dose; otherwise use age.

**DOSE** — Every 6-8 hours as needed. **DO NOT GIVE MORE THAN 4 DOSES IN 24 HOURS.**



#### Infants' MOTRIN® Concentrated Drops

**Active Ingredient:**  
Ibuprofen 50mg (NSAID)<sup>†</sup>  
(in each 1.25 mL)

<sup>†</sup> Nonsteroidal anti-inflammatory drug  
Use only as directed.

Available in:  
dye- Dye-free  
free Berry



#### Children's MOTRIN® Oral Suspension

**Active Ingredient:**  
Ibuprofen 100mg (NSAID)<sup>†</sup>  
(in each 5 mL or 1 tsp)

<sup>†</sup> Nonsteroidal anti-inflammatory drug  
Use only as directed.

Available in:  
 Original  
Berry  
dye- Dye-free  
free Berry

Weight (lbs)	Age		
<b>6-11 lbs</b>	0-5 mos	<b>Do not use</b>	—
<b>12-17 lbs</b>	6-11 mos	<b>1.25 mL</b>	
<b>18-23 lbs</b>	12-23 mos	<b>1.875 mL</b>	
<b>24-35 lbs</b>	2-3 yrs	—	<b>5 mL (1 tsp)</b>
<b>36-47 lbs</b>	4-5 yrs	—	<b>7.5 mL (1 ½ tsp)</b>
<b>48-59 lbs</b>	6-8 yrs	—	<b>10 mL (2 tsp)</b>
<b>60-71 lbs</b>	9-10 yrs	—	<b>12.5 mL (2 ½ tsp)</b>
<b>72-95 lbs</b>	11 yrs	—	<b>15 mL (3 tsp)</b>

### IMPORTANT INSTRUCTIONS FOR PROPER USE

- Read and follow the label on all MOTRIN® products
- Take every 6-8 hours as needed
- Do NOT exceed more than 4 doses in 24 hours
- Do NOT administer longer than 10 days, unless directed by a doctor
- Ask a doctor or pharmacist before use if the child is taking any other drug containing an NSAID<sup>†</sup> (prescription or non-prescription)
  - MOTRIN® contains ibuprofen
- Use only the dosing device that comes with a specific product

<sup>†</sup> Nonsteroidal anti-inflammatory drug

